



HAL
open science

Dynamics of a chain of permanent magnets

Joosung Lee, Jean Boisson, Corinne Rouby, Olivier Doaré, Michele Ducceschi,
Laurence Bodelot

► **To cite this version:**

Joosung Lee, Jean Boisson, Corinne Rouby, Olivier Doaré, Michele Ducceschi, et al.. Dynamics of a chain of permanent magnets. ESMC 2015 - The 9th European Solid Mechanics Conference, Jul 2015, Madrid, Spain. hal-01411963

HAL Id: hal-01411963

<https://ensta-paris.hal.science/hal-01411963v1>

Submitted on 8 Dec 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Dynamics of a Chain of Permanent Magnets

J. LEE[†], J. BOISSON[†], C. ROUBY[†], O. DOARÉ[†], M. DUCCESCHI[†], L. BODELOT[‡]

[†] Unité de Mécanique (UME), ENSTA-Paristech,
828 Boulevard des Maréchaux, 91762 Palaiseau cedex, France
joosung.lee@ensta-paristech.fr

[‡] Laboratoire de Mécanique des Solides (LMS), École Polytechnique,
Route de Saclay, 91128 Palaiseau cedex, France

ABSTRACT

An arrangement of several spherical or cylindrical magnets presents different stable configurations [1]. One of them is the straight chain (figures 1a, 1b), whose dynamics is studied in the present work. This structure behaves similarly to a beam, but here the rigidity is exclusively due to magnetic forces.




Figure 1: Sketch (a) and binary image (b) of a chain of cylindrical magnets at equilibrium and deformed configurations

Theoretically, the dynamical equations of the structure are obtained by first providing an expression of the energies involved in the system. At this stage, the magnetic interactions are either modeled by considering dipoles approximations for the magnets or by computing numerically the magnetic field around the magnets. Conditions of contact are introduced in the model thanks to Lagrange multipliers and a dynamical system governing the displacement of each magnet is finally obtained. An analogy with the eigenfrequencies of elastic beams allows to provide a model of an equivalent flexural rigidity induced by magnetic forces. Good agreement is found with the equivalent rigidity obtained for circular rings of magnets [2].

An experimental study of rigid assembly composed with neodymium permanent magnets is also performed. Free oscillations and forced oscillations experiments are realized. A good agreement is found between experimental and theoretical eigenfrequencies and eigenmodes (figure 2).

Next, the effect of an external magnetic field on the dynamics of a clamped-free chain of cylindrical magnets is studied. Here, the magnetic field is even able to modify the stability properties of the system. The clamped-free chain can now buckle in the same way as a beam

would buckle when submitted to gravity. The comparison between theoretical and experimental results emphasizes the limitations of the dipolar description for cylindrical magnets.

Using full computations of the magnetic field, we now develop dynamical models of other systems involving different shapes for the magnets (spheres, cylinders, plates...), as well as corresponding experiments.




Figure 2: Comparison of experimental and theoretical eigenfrequencies. Symbols present experimental results for different cases, plain lines show the two first theoretical eigenfrequencies, dashed and dash-dotted lines are the asymptotic behavior of theoretical eigenfrequencies for large n .

References

- [1] R. Messina and L. Abou Khalil *Physical Review E*, 89, 011202(R) (2014)
- [2] D. Vella, E. Du Pontavice, C.L. Hall and A. Goriely, *Proc. Roy. Soc. A*, 470, 20130609 (2014)